

Goat anti-Mouse IgG (H+L) Secondary Antibody, HRP

Product Details

Size	2 mL
Species Reactivity	Mouse
Host/Isotype	Goat / IgG
Class	Polyclonal
Type	Secondary Antibody
Conjugate	HRP
Form	Lyophilized
Concentration	0.8 mg/mL
Purification	Antigen affinity chromatography
Storage buffer	PBS, pH 7.6, with 15mg/mL BSA, 50mM sucrose
Contains	no preservative
Storage conditions	4°C
RRID	AB_228307

Applications	Tested Dilution	Publications
Western Blot (WB)	1:5,000-1:200,000	0 Publication
Immunohistochemistry (IHC)	-	0 Publication
Immunohistochemistry (Paraffin) (IHC (P))	Assay-dependent	0 Publication
Immunocytochemistry (ICC/IF)	-	0 Publication
ELISA (ELISA)	1:10,000-1:25,000	0 Publication
Immunoprecipitation (IP)	1:500-1:5,000	0 Publication
Miscellaneous PubMed (Misc)	-	0 Publication

Product Specific Information

Product # 31430 has been successfully used in Western blot, IHC, ELISA and IP applications.

Product # 31430 reacts with the heavy chains of mouse igg and with the light chains common to most mouse immunoglobulins, but does not react against non-immunoglobulin serum proteins. However, this antibody may cross-react with immunoglobulins from other species.


Store product at 4°C until opened. To extend the shelf-life of this product, add an equal volume of glycerol to make a final concentration of approximately 50% glycerol and store at -20°C.

Reconstitute with 2.0 mL of distilled water (0.8 mg/mL after restoration).

Product Images For Goat anti-Mouse IgG (H+L) Secondary Antibody, HRP


Mouse IgG (H+L) Secondary Antibody (31430) in WB

Chemiluminiscent western blot was performed using Goat anti-Mouse IgG (Heavy Chain) Secondary Antibody, HRP (Product # 31430). Whole cell extracts of U-87 MG (Lane 1, 2, 3, 4), SH-SY5Y (Lane 5, 6, 7, 8), HeLa (Lane 9) and Hep G2 (Lane 10) were electrophoresed using NuPAGE™ 4-12% Bis-Tris Protein Gel (Product # NP03221BOX). Resolved proteins were transferred onto anitrocellulose membrane (Product # IB23001) by iBlot® 2 Dry BlottingSystem (Product # IB21001). The blot was probed with PGP9.5 Monoclonal Antibody (13C4) (Product # MA1-90008), and HSP70 Polyclonal Antibody (Product # PA5-28003). Secondary antibodies (Product # 31430, 1:30,000), and (Product # A27036, 1:20,000) were used for detection of PGP9.5, and HSP70 respectively. Chemiluminiscent detection was performed using iBright™FL1500 (Product # A44115). The anti-mouse secondary antibody (Product # 31430) specifically detects the mouse primary antibody.




Mouse IgG (H+L) Secondary Antibody (31430) in WB

Western blot analysis was performed on whole cell extracts (30 µg lysate) of K-562 (Lane 1) and U87-MG (Lane 2). The blots were probed with Anti-SOD2 Mouse Monoclonal Antibody (Product # MA1-106, 0.25 µg/mL) and detected by chemiluminescence using Goat anti-Mouse IgG (H+L) Secondary Antibody, HRP conjugate (Product # 31430) at dilutions 1:5,000 (Fig. 1), 1:100,000 (Fig. 2) and 1: 200,000 (Fig. 3). A 25 kDa band corresponding to SOD2 was observed. Known quantity of protein samples were electrophoresed using Novex® NuPAGE® 12 % Bis-Tris gel (Product # NP0342BOX), XCell SureLock™ Electrophoresis System (Product # EI0002) and Novex® Sharp Pre-Stained Protein Standard (Product # LC5800). Resolved proteins were then transferred onto a nitrocellulose membrane with iBlot® 2 Dry Blotting System (Product # IB21001). The membrane was probed with the relevant primary and secondary antibody after blocking with 5 % skimmed milk. Chemiluminescent detection was performed using Pierce™ ECL Western Blotting Substrate (Product # 32106).



Mouse IgG (H+L) Secondary Antibody (31430) in WB

Western blot analysis of beta-3 Tubulin was performed by loading 25 µg of various brain, SHSY5Y, and HeLa cell lysates and 10 µL of PageRuler Prestained Protein Ladder (Product # 26616) onto a 4-20% Tris-HCl polyacrylamide gel. Proteins were transferred to a PVDF membrane and blocked with 5% BSA/TBST for at least 1 hour. The membrane was probed with a beta-3 Tubulin monoclonal antibody (Product # MA1-118) at a dilution of 1:2000 overnight at 4°C on a rocking platform, washed in TBS-0.1%Tween-20, and probed with a goat anti-mouse IgG-HRP secondary antibody (Product # 31430) at a dilution of 1:20,000 for 1 hour. Chemiluminescent detection was performed using SuperSignal West Pico (Product # 34080). Note: analysis indicates both reactive (neuronal) and non-reactive negative control (HeLa cell lysate) for beta-3 Tubulin using Product # MA1-118.



View more figures on thermofisher.com

Hydrophobin-Fused SARS-CoV-2 RBD production in *Nicotiana benthamiana* L. for COVID-19 serology. *Future Sci OA* (2025)

Risk assessment of 2024 cattle H5N1 using age-stratified serosurveillance data. *Emerg Microbes Infect* (2025)

Histone acetyltransferase 1 promotes ovarian cancer progression by regulating cell proliferation and the cell cycle. *Oncol Lett* (2025)

A comparative proteomic, transcriptomic and glycomic analysis of extracellular vesicle isolation techniques highlights ExoGAG efficiency for a more complete identification of breast milk molecular signaling pathways. *Cell Commun Signal* (2025)

Genome-wide mapping of arsenic-activated Nrf2 reveals metabolic and epigenetic reprogramming in induced pluripotent stem cells. *Redox Biol* (2025)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.