

PRODUCT INFORMATION

TopVision Agarose Tablets

| C | g | X | 1000 | lak |
|---|---|----------|-------|------------|
| | C | ၁ | b g x | 5 g x 1000 |

| Lot: _ | Exp |). | |
|--------|-----|----|--|
| | | | |

Sizes available:

| Cat No | Description | Pack Size |
|--------|---------------------------|------------------|
| #R2801 | TopVision Agarose Tablets | 0.5 g × 200 tab |
| #R2802 | TopVision Agarose Tablets | 0.5 g × 1000 tab |

Country of Origin: China Assembled in (EU) Lithuania

Store at room temperature, away from light and moisture

Description:

TopVision Agarose Tablets is a low-electro endosmosis, multi-purpose agarose in tablet form, delivered in a convenient blister pack. It is made of standard melting point agarose, which yields high resolution sharp DNA bands with high clarity and low background. Its optimized gel strength enhances ease of gel processing and handling.

Unlike traditional agarose powder that relies heavily on the use of toxic organic solvents, TopVision Agarose Tablets are manufactured using an innovative Organic Solvent-Free Manufacturing process that is more environmentally friendly.

Specifications:

| opeomedions. | |
|---|---|
| CAS | 9012-36-6 |
| Appearance | White to off-white powder |
| EEO (electroendosmosis) | ≤ 0.13 |
| Gelling Point | 36 °C ± 1.5 °C (1.5% gel) |
| Melting Point | 88 °C ± 1.5 °C (1.5% gel) |
| Solubility | Clear colorless solution at 1 g in 100 mL water |
| Moisture | ≤ 10% |
| Gel Strength | ≥ 1200 g/cm² (1% Gel) |
| Sulfate | ≤ 0.15% |
| Ash | ≤ 0.5% |
| Impurities (such as DNase, RNase, Protease, and Endonuclease) | None |

Protocol:

1. Add appropriate number of agarose tablets to the electrophoresis buffer based on the table below to prepare your desired gel percentage.

Note: Use a flask that is 2 to 4 times the volume of the solution being prepared.

| Gel % | 1 tablet | 2 tablets | 3 tablets |
|-------|----------|-----------|-----------|
| 1% | 50 mL | 100 mL | 150 mL |
| 1.2% | 42 mL | 83 mL | 125 mL |
| 1.3% | 38 mL | 77 mL | 115 mL |
| 1.5% | 33 mL | 67 mL | 100 mL |
| 1.8% | 28 mL | 56 mL | 83 mL |
| 2% | 25 mL | 50 mL | 75 mL |

- 2. Before heating soak tablets in a buffer (~ 4 minutes) until tablets completely break into fine-particle slurry. Swirl the slurry to break up any remaining particles.

 Important: Ensure tablets break up entirely. Heating will render non-dispersed agarose particles insoluble.
- 3. Heat the slurry in a microwave on high power setting until it starts to boil, allow boiling for ~ 30 seconds.
- 4. Note: Heating times are dependent on the volume of liquid and number of gel tablets to dissolve.
- 5. Remove the flask from microwave, swirl gently to dissolve any remaining agarose particles.

- 6. Reheat on high power for 1-2 minutes or until the solution is clear and all particles are dissolved.
- 7. Remove the flask from the microwave oven, and gently swirl.
- 8. Cool the solution to approximately 60 °C before pouring.
- 9. Pour the gel.

If a boiling water bath is used:

To melt agarose, simply heat the slurry in a boiling water bath. Bring the solution to a boil and allow it to boil for 5-10 minutes stirring continuously until agarose dissolves completely.

Note: Use caution when handling as solution may be extremely hot.

Recommended agarose percentage for DNA separation

| Agarose concentration | 0.3% | 0.6% | 0.7% | 0.9% | 1.2% | 1.5% | 2.0% |
|-----------------------|------|------|--------|-------|-------|-------|-------|
| DNA size range (kb) | 5-60 | 1-20 | 0.8-10 | 0.5-7 | 0.4-6 | 0.2-4 | 0.1-3 |

Quality control: all parameters stated in the specifications are tested and confirmed.

LIMITED USE LABEL LICENSE: Internal Research and Development Use Only.

The purchase of this product conveys to the buyer the limited, non-exclusive, non-transferable right (without the right to resell, repackage, or further sublicense) to use this product for internal research and development purposes. No other license is granted to the buyer whether expressly, by implication, by estoppel or otherwise. In particular, the purchase of the product does not include or carry any right or license to use, develop, or otherwise exploit this product commercially and no rights are conveyed to the buyer to use the product or components of the product for purposes including but not limited to provision of services to a third party, generation of commercial databases or clinical diagnostics. This product is sold pursuant to authorization from Thermo Fisher Scientific and Thermo Fisher Scientific reserves all other rights. For information on purchasing a license for uses other than internal research and development purposes, please contact outlicensing@lifetech.com or Out Licensing, Life Technologies Inc., 5791 Van Allen Way, Carlsbad, California 92008.

PRODUCT USE LIMITATION

This product is developed, designed and sold exclusively *for research purposes and in vitro use only.* The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals. Please refer to www.thermoscientific.com/onebio for Material Safety Data Sheet of the product.

© 2015 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.