

# Human M-CSF Recombinant Protein

## Product Details

Size	10 µg
Species	Human
Published Species	Human, Mouse
Expression system	HEK293 cells
Molecular weight	38 kDa
Class	Recombinant
Type	Protein
Purity	>95% by SDS-PAGE
Endotoxin concentration	<0.1 ng/µg
Activity	ED50 5 ng/mL; determined by the dose-dependent proliferation of M-NSF60 cells.
Conjugate	Unconjugated
Form	Lyophilized
Purification	purified
Contains	no preservative
Storage conditions	4° C

Applications	Tested Dilution	Publications
Western Blot (WB)	Assay-dependent	-
Immunohistochemistry (IHC)	Assay-dependent	-
ELISA (ELISA)	Assay-dependent	-
Functional Assay (Functional)	Assay-dependent	5 Publications
Control (Ctrl)	Assay-dependent	-

## Product Specific Information

**Reconstitution:** Lyophilized human M-CSF should be reconstituted in 20mM Tris-HCl, pH 8.0 to a concentration of 0.1-1.0 mg/mL. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein, such as 0.1% BSA and store in working aliquots at -20°C to -80°C.

**Storage:** Lyophilized human M-CSF should be stored at 2°C to 8°C, preferably desiccated. Store reconstituted human M-CSF at -20°C (not in a frost-free freezer). Keep freeze-thaw cycles to a minimum.

## Functional Assay (5)

Molecular therapy. Nucleic acids

### FABP4 facilitates inflammasome activation to induce the Treg/Th17 imbalance in preeclampsia via forming a positive feedback with IL-17A.

Authors: Chang GP,Yang XL,Liu W,Lin S,Yang SL,Zhao MY

Year  
2021

Species  
Human

Dilution  
50 ng/mL

Stem cell reports

### Neuroinflammation and EIF2 Signaling Persist despite Antiretroviral Treatment in an hiPSC Tri-culture Model of HIV Infection.

Authors: Ryan SK,Gonzalez MV,Garifallou JP,Bennett FC,Williams KS,Sotuyo NP,Mironets E,Cook K,Hakonarson H,Anderson SA,Jordan-Sciutto KL

Year  
2020

Species  
Human

Dilution  
25 ng/mL

[View more Functional references on thermofisher.com](#)

## More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON-INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.