



IL-10 Monoclonal Antibody (JES3-9D7), PerCP-eFluor™ 710, eBioscience™

Product Details	
Size	25 Tests
Species Reactivity	Human
Published Species	Human
Host/Isotype	Rat / IgG1, kappa
Recommended Isotype Control	Rat IgG1 kappa Isotype Control (eBRG1), PerCP-eFluor™ 710, eBioscience™
Class	Monoclonal
Туре	Antibody
Clone	JES3-9D7
Conjugate	PerCP-eFluor™ 710
Excitation/Emission Max	482/708 nm
Form	Liquid
Concentration	5 μL/Test
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2, with 0.2% BSA
Contains	0.09% sodium azide
Storage conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_2573832

Applications	Tested Dilution	Publications
Flow Cytometry (Flow)	5 μL (0.03 μg)/test	9 Publications
ELISA (ELISA)	-	1 Publication
Functional Assay (Functional)	-	1 Publication

Product Specific Information

Description: The JES3-9D7 monoclonal antibody reacts with human interleukin-10 (IL-10).

Applications Reported: This JES3-9D7 antibody has been reported for use in intracellular staining followed by flow cytometric analysis.

Applications Tested: This JES3-9D7 antibody has been pre-titrated and tested by intracellular staining and flow cytometric analysis of stimulated normal human peripheral blood cells using the Intracellular Fixation & Permeabilization Bufer Set (Product # 88-8824-00) and protocol. Please refer to Best Protocols: Protocol A: Two step protocol for (cytoplasmic) intracellular proteins. This can be used at 5 μ L (0.03 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells /test.

PerCP-eFluor® 710 emits at 710 nm and is excited with the blue laser (488 nm); it can be used in place of PerCP-Cyanine5.5. We recommend using a 710/50 bandpass filter, however, the 695/40 bandpass filter is an acceptable alternative. Please make sure that your instrument is capable of detecting this fluorochrome.


Light sensitivity: This tandem dye is sensitive to photo-induced oxidation. Please protect this vial and stained samples from light.

Fixation: Samples can be stored in IC Fixation Buffer (Product # 00-8222) (100 μ L of cell sample + 100 μ L of IC Fixation Buffer) or 1-step Fix/Lyse Solution (Product # 00-5333) for up to 3 days in the dark at 4°C with minimal impact on brightness and FRET efficiency/compensation. Some generalizations regarding fluorophore performance after fixation can be made, but clone specific performance should be determined empirically.

Excitation: 488 nm; Emission: 710 nm; Laser: Blue Laser.

Filtration: 0.2 µm post-manufacturing filtered.

Product Images For IL-10 Monoclonal Antibody (JES3-9D7), PerCP-eFluor™ 710, eBioscience™



IL-10 Antibody (46-7108-41) in Flow

Normal human peripheral blood mononuclear cells were cultured for 6 days under Th2-polarizing conditions. The cells were unstimulated (left) or restimulated (right) with the Cell Stimulation Cocktail (plus protein transport inhibitors) (Product # 00-4975-03) then intracellularly stained with Anti-Human CD4 FITC (Product # 11-0049-42) and Anti-Human IL-10 PerCP-eFluor® 710 using the Intracellular Fixation & Permeabilization Buffer Set (Product # 88-8824-00) and protocol. Total viable cells, as determined by Fixable Viability Dye eFluor® 450 (Product # 65-0863-14), were used for analysis.

View more figures on thermofisher.com

□ 11 References

Flow Cytometry (9)

Journal of cardiovascular development and disease

Circulating Regulatory B-Lymphocytes in Patients with Acute Myocardial Infarction: A Pilot Study.

"Published figure using IL-10 monoclonal antibody (Product # 46-7108-42) in Flow Cytometry"

Authors: Volodarsky I,Shimoni S,Haberman D,Mirkin V,Fabrikant Y,Yoskovich Mashriki T,Zalik A,George J

Year 2022

Journal of inflammation research

Altered T-Cell Subsets are Associated with Dysregulated Cytokine Secretion of CD4⁺ T Cells During HIV Infection.

"Published figure using IL-10 monoclonal antibody (Product # 46-7108-42) in Flow Cytometry" Authors: Wang D,Jiang Y,Song Y,Zeng Y,Li C,Wang X,Liu Y,Xiao J,Kong Y,Zhao H

Year 2022

View more Flow references on thermofisher.com

ELISA (1)

Cellular immunology

Granzyme B production distinguishes recently activated CD8(+) memory cells from resting memory cells.

Authors: Nowacki TM, Kuerten S, Zhang W, Shive CL, Kreher CR, Boehm BO, Lehmann PV, Tary-Lehmann M

Year

Functional Assay (1)

Journal of immunology (Baltimore, Md.: 1950)

Mycobacterium bovis BCG attenuates surface expression of mature class II molecules through IL-10-dependent inhibition of cathepsin S.

Authors: Sendide K,Deghmane AE,Pechkovsky D,Av-Gay Y,Talal A,Hmama Z

Year 2005

More applications with references on thermofisher.com

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization, Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production of specifications and/or accompanying package inserts ("Documentation"), No claim of situability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, in kis warranty and in the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED. OR PEPAIR, REPLACE OR REFUND FOR RIFE NON-CONFORMING PRODUCTS, AT SELLEYS SOLE OPTION. THERE IS NO GBLIGATION TO REPAIR, REPLACE OR REFUND FOR RIFE NON-CONFORMING PRODUCTS, AT SELLEYS SOLE OPTION. THERE IS NO GBLIGATION TO REPAIR, REPLACE OR REFUND FOR RIFE NON-CONFORMING PRODUCTS, AT SELLEY SOLE OPTION. THERE IS NO GBLIGATION TO REPAIR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS AT THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or vivo or in vivo therapeutic uses, or any type of consumption by or application to human or a