

PRODUCT INFORMATION

Eco31I (Bsal)

#ER0292 5000 U

Lot: ____ **Expiry Date:** _

5'...G G T C T C (N) $_1$ \dagger...3' 3'...C C A G A G (N)_{5↑}...5'

Concentration: 10 U/μL

F.coli that carries the cloned eco31IR Source:

aene from E.coli RFL31

2x1 mL of 10X Buffer G Supplied with:

1 mL of 10X Buffer Tango

Store at -20°C













In total 4 vials.

BSA included

RECOMMENDATIONS

1X Buffer G (for 100% Eco31l digestion) 10 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, 50 mM NaCl, 0.1 mg/mL BSA.

Incubation temperature

37°C.

Unit Definition

One unit is defined as the amount of Eco311 required to digest 1 µg of lambda DNA dcm⁻-Hindlll fragments in 1 hour at 37°C in 50 µL of recommended reaction buffer. Assayed using lambda DNA dcm⁻, as one of two Eco311 recognition sites is difficult to cleave.

Dilution

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

Double Digests

Thermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango[™] Buffer. Please refer to www.thermoscientific.com/doubledigest to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/mL BSA.

Rev.12

Storage Buffer

Eco31I is supplied in: 10 mM Tris-HCI (pH 7.5 at 25°C), 200 mM KCI, 1 mM DTT, 0.1 mM EDTA, 0.2 mg/mL BSA and 50% glycerol.

Recommended Protocol for Digestion

• Add:

nuclease-free water 16 μ L 10X Buffer G 2 μ L DNA (0.5-1 μ g/ μ L) 1 μ L Eco31I 0.5-2 μ L*

- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours*.

Recommended Protocol for Digestion of PCR Products Directly after Amplification

• Add:

PCR reaction mixture $10 \mu L$ (about 1 μg of DNA)

Water, nuclease-free $16 \mu L$ 10X Buffer G $2 \mu L$ Eco31I $1-2 \mu L*$

- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours*.

Thermal Inactivation

Eco31I is inactivated by incubation at 65°C for 20 min.

ENZYME PROPERTIES

Enzyme Activity in Thermo Scientific REase Buffers, % B G O R Tango 2X Tango

В	G	0	R	Tango	2X Tango
50-100	100	0-20	0-20	50-100	20-50

Methylation Effects on Digestion

Dam: never overlaps – no effect.

Dcm: may overlap — cleavage impaired. CpG: may overlap — cleavage impaired.

EcoKI: never overlaps – no effect.

EcoBl: may overlap – effect not determined.

Stability during Prolonged Incubation

A minimum of 0.3 units of the enzyme is required for complete digestion of 1 μ g of lambda DNA dcm^- in 16 hours at 37°C.

Digestion of Agarose-embedded DNA

A minimum of 5 units of the enzyme is required for complete digestion of 1 μ g of agarose-embedded lambda DNA dcm^- in 16 hours.

Number of Recognition Sites in DNA

λ	ФХ174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
2	0	1	1	1	1	0

Note

Eco31I cleavage is impaired by overlapping *dcm* methylation. To avoid *dcm* methylation, use a *dam*⁻, *dcm*⁻ strain such as GM2163 (#M0099).

For **CERTIFICATE OF ANALYSIS** see back page

See Overdigestion Assay.

CERTIFICATE OF ANALYSIS

Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 80-fold overdigestion with Eco31I (5 U/ μ g lambda DNA $dcm^- \times 16$ hours).

Ligation and Recleavage (L/R) Assay

The ligation and recleavage assay was replaced with LO test after validating experiments showed LO test ability to trace nuclease and phosphatase activities with sensitivity that is higher than L/R by a factor of 100.

Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or doublestranded labeled oligonucleotides occurred during incubation with 10 units of Eco31I for 4 hours.

Quality authorized by:



Jurgita Zilinskiene

PRODUCT USE LIMITATION

This product is developed, designed and sold exclusively *for research purposes and in vitro use only.* The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.thermoscientific.com/onebio for Material Safety Data Sheet of the product.

© 2012 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.